

**September 2017 *Journal of Environmental Health***  
**Volume 80, Number 2**

**Direct From ATSDR: Community Exposure to Per- and Polyfluoroalkyl Substances in Drinking Water: A National Issue**

References

- Alexander, B.H., & Olsen, G.W. (2007). Bladder cancer in perfluorooctanesulfonyl fluoride manufacturing workers. *Annals of Epidemiology*, *17*(6), 471–478.
- Barry, V., Winquist, A., & Steenland, K. (2013). Perfluorooctanoic acid (PFOA) exposures and incident cancers among adults living near a chemical plant. *Environmental Health Perspectives*, *121*(11–12), 1313–1318.
- Bellinger, D.C. (2013). Prenatal exposures to environmental chemicals and children's neurodevelopment: An update. *Safety and Health at Work*, *4*(1), 1–11.
- Butenhoff, J., Costa, G., Elcombe, C., Farrar, D., Hansen, K., Iwai, H., . . . Thomford, P. (2002). Toxicity of ammonium perfluorooctanoate in male cynomolgus monkeys after oral dosing for 6 months. *Toxicological Sciences*, *69*(1), 244–257.
- Butenhoff, J.L., Chang, S.C., Olsen, G.W., & Thomford, P.J. (2012). Chronic dietary toxicity and carcinogenicity study with potassium perfluorooctanesulfonate in Sprague Dawley rats. *Toxicology*, *293*(1–3), 1–15.
- Centers for Disease Control and Prevention. (2017). *National report on human exposure to environmental chemicals: Updated tables, January 2017*. Retrieved from <https://www.cdc.gov/exposurereport>
- Chen, M.H., Ha, E.H., Wen, T.W., Su, Y.N., Lien, G.W., Chen, C.Y., . . . Hsieh, W.S. (2012). Perfluorinated compounds in umbilical cord blood and adverse birth outcomes. *PLoS One*, *7*(8), e42474.
- Crawford, N.M., Fenton, S.E., Strynar, M., Hines, E.P., Pritchard, D.A., & Steiner, A.Z. (2017). Effects of perfluorinated chemicals on thyroid function, markers of ovarian reserve, and natural fertility. *Reproductive Toxicology*, *69*, 53–59.
- Cui, L., Zhou, Q.F., Liao, C.Y., Fu, J.J., & Jiang, G.B. (2009). Studies on the toxicological effects of PFOA and PFOS on rats using histological observation and chemical analysis. *Archives of Environmental Contamination and Toxicology*, *56*(2), 338–349.
- Dalsager, L., Christensen, N., Husby, S., Kyhl, H., Nielsen, F., Høst, A., . . . Jensen, T.K. (2016). Association between prenatal exposure to perfluorinated compounds and symptoms of infections at age 1–4 years among 359 children in the Odense Child Cohort. *Environment International*, *96*, 58–64.
- Darrow, L.A., Stein, C.R., & Steenland, K. (2013). Serum perfluorooctanoic acid and perfluorooctane sulfonate concentrations in relation to birth outcomes in the mid-Ohio valley, 2005–2010. *Environmental Health Perspectives*, *121*(10), 1207–1213.
- DeWitt, J.C. (2015). *Toxicological effects of perfluoroalkyl and polyfluoroalkyl substances*. Switzerland: Springer International Publishing.
- Eriksen, K.T., Raaschou-Nielsen, O., McLaughlin, J.K., Lipworth, L., Tjønneland, A., Overvad, K., & Sørensen, M. (2013). Association between plasma PFOA and PFOS levels and total cholesterol in a middle-aged Danish population. *PLoS One*, *8*(2), e56969.
- Fei, C., McLaughlin, J.K., Lipworth, L., & Olsen, J. (2009). Maternal levels of perfluorinated chemicals and subfecundity. *Human Reproduction*, *24*(5), 1200–1205.

- Fei, C., McLaughlin, J.K., Tarone, R.E., & Olsen, J. (2007). Perfluorinated chemicals and fetal growth: A study within the Danish National Birth Cohort. *Environmental Health Perspectives*, 115(11), 1677–1682.
- Fei, C., & Olsen, J. (2011). Prenatal exposure to perfluorinated chemicals and behavioral or coordination problems at age 7 years. *Environmental Health Perspectives*, 119(4), 573–578.
- Fitz-Simon, N., Fletcher, T., Luster, M.I., Steenland, K., Calafat, A.M., Kato, K., & Armstrong, B. (2013). Reductions in serum lipids with a 4-year decline in serum perfluorooctanoic acid and perfluorooctanesulfonic acid. *Epidemiology*, 24(4), 569–576.
- Fletcher, T., Galloway, T.S., Melzer, D., Holcroft, P., Cipelli, R., Pilling, L.C., . . . Harries, L.W. (2013). Associations between PFOA, PFOS and changes in the expression of genes involved in cholesterol metabolism in humans. *Environment International*, 57–58: 2–10.
- Forns, J., Iszatt, N., White, R.A., Mandal, S., Sabaredzovic, A., Lamoree, M., . . . Eggesbø, M. (2015). Perfluoroalkyl substances measured in breast milk and child neuropsychological development in a Norwegian birth cohort study. *Environment International*, 83, 176–182.
- Frisbee, S.J., Shankar, A., Knox, S.S., Steenland, K., Savitz, D.A., Fletcher, T., & Ducatman, A.M. (2010). Perfluorooctanoic acid, perfluorooctanesulfonate, and serum lipids in children and adolescents: results from the C8 Health Project. *Archives of Pediatric & Adolescent Medicine*, 164(9), 860–869.
- Grandjean, P., Andersen, E.W., Budtz-Jørgensen, E., Nielsen, F., Mølbak, K., Weihe, P., & Heilmann, C. (2012). Serum vaccine antibody concentrations in children exposed to perfluorinated compounds. *JAMA*, 307(4), 391–397.
- Granum, B., Haug, L.S., Namork, E., Stolevik, S.B., Thomsen, C., Aaberge, I.S., . . . Nygaard, U.C. (2013). Pre-natal exposure to perfluoroalkyl substances may be associated with altered vaccine antibody levels and immune-related health outcomes in early childhood. *Journal of Immunotoxicology*, 10(4), 373–379.
- Iwai, H., & Yamashita, K. (2006). A fourteen-day repeated dose oral toxicity study of APFO in rats. *Drug and Chemical Toxicology*, 29(3), 323–332.
- Jain, R.B. (2013). Association between thyroid profile and perfluoroalkyl acids: Data from NHNAES 2007–2008. *Environmental Research*, 126, 51–59.
- Joensen, U.N., Veyrand, B., Antignac, J.P., Blomberg Jensen, M., Petersen, J.H., Marchand, P., . . . Jørgensen, N. (2013). PFOS (perfluorooctanesulfonate) in serum is negatively associated with testosterone levels, but not with semen quality, in healthy men. *Human Reproduction*, 28(3), 599–608.
- Lau, C., Butenhoff, J.L., & Rogers, J.M. (2004). The developmental toxicity of perfluoroalkyl acids and their derivatives. *Toxicology and Applied Pharmacology*, 198(2), 231–241.
- Lau, C., Thibodeaux, J.R., Hanson, R.G., Narotsky, M.G., Rogers, J.M., Lindstrom, A.B., & Strynar, M.J. (2006). Effects of perfluorooctanoic acid exposure during pregnancy in the mouse. *Toxicological Sciences*, 90(2), 510–518.
- Lopez-Espinosa, M.-J., Mondal, D., Armstrong, B., Bloom, M.S., & Fletcher, T. (2012). Thyroid function and perfluoroalkyl acids in children living near a chemical plant. *Environmental Health Perspectives*, 120(7), 1036–1041.
- Luebker, D.J., York, R.G., Hansen, K.J., Moore, J.A., & Butenhoff, J.L. (2005). Neonatal mortality from in utero exposure to perfluorooctanesulfonate (PFOS) in Sprague-Dawley rats: Dose-response, and biochemical and pharmacokinetic parameters. *Toxicology*, 215(1–2), 149–169.

- Mastrantonio, M., Bai, E., Uccelli, R., Cordiano, V., Screpanti, A., & Crosignani, P. (2017). Drinking water contamination from perfluoroalkyl substances (PFAS): an ecological mortality study in the Veneto Region, Italy. *European Journal of Public Health*. Advanced online publication.
- Nelson, J.W., Hatch, E.E., & Webster, T.F. (2010). Exposure to polyfluoroalkyl chemicals and cholesterol, body weight, and insulin resistance in the general U.S. population. *Environmental Health Perspectives*, *118*(2), 197–202.
- Sakr, C.J., Kreckmann, K.H., Green, J.W., Gillies, P.J., Reynolds, J.L., & Leonard, R.C. (2007). Cross-sectional study of lipids and liver enzymes related to a serum biomarker of exposure (ammonium perfluorooctanoate or APFO) as part of a general health survey in a cohort of occupationally exposed workers. *Journal of Occupational and Environmental Medicine*, *49*(10), 1086–1096.
- Seacat, A.M., Thomford, P.J., Hansen, K.J., Olsen, G.W., Case, M.T., & Butenhoff, J.L. (2002). Subchronic toxicity studies on perfluorooctanesulfonate potassium salt in cynomolgus monkeys. *Toxicological Sciences*, *68*(1), 249–264.
- Shrestha, S., Bloom, M.S., Yucel, R., Seegal, R.F., Rej, R., McCaffrey, R.J., . . . Fitzgerald, E.F. (2016). Perfluoroalkyl substances, thyroid hormones, and neuropsychological status in older adults. *International Journal of Hygiene and Environmental Health*, *220*(4), 647–782.
- Steenland, K., Tinker, S., Frisbee, S., Ducatman, A., & Vaccarino, V. (2009). Association of perfluorooctanoic acid and perfluorooctane sulfonate with serum lipids among adults living near a chemical plant. *American Journal of Epidemiology*, *170*(10), 1268–1278.
- Steenland, K., Zhao, L., Winquist, A., & Parks, C. (2013). Ulcerative colitis and perfluorooctanoic acid (PFOA) in a highly exposed population of community residents and workers in the mid-Ohio valley. *Environmental Health Perspectives*, *121*(8), 900–905.
- Stein, C. R., & Savitz, D.A. (2011). Serum perfluorinated compound concentration and attention deficit/hyperactivity disorder in children 5–18 years of age. *Environmental Health Perspectives*, *119*(10), 1466–1471.
- Stein, C. R., Savitz, D.A., & Bellinger, D.C. (2013). Perfluorooctanoate and neuropsychological outcomes in children. *Epidemiology*, *24*(4), 590–599.
- U.S. Census Bureau. (2007). *American housing survey for the United States: 2007*. Washington, DC: U.S. Government Printing Office.
- U.S. Environmental Protection Agency. (2017). *Occurrence data for the unregulated contamination monitoring rule: UCMR 3 (2012–2015) occurrence data*. Retrieved from <https://www.epa.gov/dwucmr/occurrence-data-unregulated-contaminant-monitoring-rule#3>
- Vieira, V.M., Hoffman, K., Shin, H.-M., Weinberg, J.M., Webster, T.F., & Fletcher, T. (2013). Perfluorooctanoic acid exposure and cancer outcomes in a contaminated community: a geographic analysis. *Environmental Health Perspectives*, *121*(3), 318–323.
- Wang, Y., Rogan, W.J., Chen, P.C., Lien, G.W., Chen, H.Y., Tseng, Y.C., . . . Wang, S.L. (2014). Association between maternal serum perfluoroalkyl substances during pregnancy and maternal and cord thyroid hormones: Taiwan maternal and infant cohort study. *Environmental Health Perspectives*, *122*(5), 529–534.

Whitworth, K.W., Haug, L.S., Baird, D.D., Becher, G., Hoppin, J.A., Skjaerven, R., . . . Longnecker, M.P. (2012). Perfluorinated compounds and subfecundity in pregnant women. *Epidemiology*, 23(2), 257–263.